David N. Lieberman,¹ Grace Correa,¹ Arthur Beisang,² Robin Ryther,³ Christopher W. Beatty,⁴ Emily De Los Reyes,⁴ Bernhard Suter,⁵ Alan K. Percy,⁶ Amitha L. Ananth,⁶ Caroline Buchanan,⁷ Louise Cosand,⁸ Alyssa Peckham⁸

Achievement of Trofinetide Label Dose With Titration

BACKGROUND

- Rett syndrome (RTT) is a rare neurodevelopmental disorder characterized by loss of verbal communication with limited nonverbal skills, loss of fine and gross motor function, behavioral issues, seizures, hand stereotypies, and gastrointestinal problems^{1,2}
- Trofinetide was approved by the US Food and Drug Administration in March 2023 for the treatment of RTT in adults and pediatric patients aged ≥ 2 years³
- Trofinetide is recommended to be administered orally twice daily (BID), in the morning and evening, with or without food, according to patient weight (**Table 1**)³
- In LAVENDER, the randomized, placebo-controlled, phase 3 study of trofinetide in girls and women with RTT, approximately one-third of patients in the trofinetide arm had their dose adjusted for tolerability reasons⁴ and evidence suggests that clinicians are using alternative dosing strategies in the real world⁵

Table 1. Trofinetide Weight-Based Dosing Recommendations

Patient weight, kg	Trofinetide dose (BID), mg	Trofinetide volume (BID), mL
9 – <12	5000	25
12 - <20	6000	30
20 - <35	8000	40
35 - <50	10,000	50
≥50	12,000	60

BID, twice daily

OBJECTIVE

• To explore real-world alternative dosing strategies of trofinetide in patients with RTT

METHODS

Survey Design

- An electronic prescriber experience survey was designed to collect real-world trofinetide dosing strategies, dosing considerations, and achievable dose
- The survey was sent in May 2024 to 33 prescribers at 18 US RTT centers of excellence (COEs) designated by the International Rett Syndrome Foundation

Data Analysis

• All results were summarized with descriptive statistics

RESULTS

Survey Participation

- Overall, 22 prescribers from 16 COEs completed the electronic survey
- The prescribers targeted for survey completion accounted for 38.1% of trofinetide prescriptions in the United States since approval (**Table 2**)

Table 2. Electronic Survey Participation and Representation

	US trofinetide prescribers (N = 697)ª	US trofinetide prescriptions (N = 1553) ^a
Non-COE providers, n (%)	665 (95.4)	962 (61.9)
COE providers, n (%)	32 (4.6) ^b	591 (38.1)

^aAs of May 2024. ^bOne eligible prescriber was associated with zero prescriptions COE, center of excellence





Real-World Use of Trofinetide: A Survey of Dosing Strategies From US Rett Syndrome Centers of Excellence

¹Boston Children's Hospital, Boston, MA, USA; ²Gillette Children's Specialty Healthcare, St Paul, MN, USA; ³Washington University School of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State University College of Medicine, St Louis, MO, USA; ⁴Nationwide Children's Hospital, The Ohio State Uni Columbus, OH, USA; ⁵Baylor College of Medicine, Houston, TX, USA; ⁶University of Alabama at Birmingham, AL, USA; ⁷Greenwood Genetic Center, Greenville, SC, USA; ⁸Acadia Pharmaceuticals Inc., San Diego, CA, USA

Trofinetide Dosing at Treatment Initiation

• Most survey respondents (95%, n = 21) indicated that they titrate trofinetide in treatment-naïve patients with RTT rather than initiate at label dose (**Figure 1**)

Figure 1. Survey Respondents' Initial **Trofinetide Dosing Strategy**

Survey question:

How do you initiate trofinetide in a patient that has never received trofinetide before?

 Titration protocols included initiating trofinetide at a lower percentage of label dose (50%, n = 11) or lower milliliters than label dose (27%, n = 6), or were determined based on baseline conditions, such as diarrhea, constipation, and dose goal (14%, n = 3), or are unknown (9%, n = 2) (**Table 3**)

Most titration protocols start at 25–50% of the dose goal BID and increase by 5–10 mL BID every 1 to 2 weeks as tolerated

Slower titration protocols and titration protocols with lower dose increases were reported for patients with lower weight, younger patients, patients with a history of diarrhea, patients with history of poor trofinetide tolerability, and patients who experienced medication side effects

 Faster titration protocols and titration protocols with higher dose increases were reported for patients with higher weight, older patients, patients with a history of constipation, and patients with history of acceptable trofinetide tolerability

Table 3. Trofinetide Titration Protocols Reported by Survey Respondents

Titration protocols by percentage of label dose (n = 11)				
Starting dose (BID)		Titration		
25%		Increase by 25% every 2 weeks		
25%		Increase by 5 mL BID every 1 to 2 week		
33%		Increase by 33% weekly		
25–33%		Increase by 5 mL weekly		
25–50%		Increase by 5 mL BID weekly		
50%		Increase by 25% weekly		
50%		Increase by 25% weekly		
50%		Increase by 5 mL BID weekly		
50%		Increase by 5 mL BID every 1 to 2 week		
50%		Increase by 5 mL BID every 2 weeks		
50%		Increase by 5–10 mL weekly		
Titration protocols by mL of label dose (n = 6)				
Starting dose (BID)		Titration		
5 mL		Increase by 2.5 mL BID weekly		
5 mL		Increase by 5 mL weekly		
5 mL		Increase by 5 mL weekly		
5 mL		Increase by 5 mL BID weekly		
5 mL		Increase by 5 mL BID weekly		
10–15 mL		Increase by 5 mL BID weekly		
Titration protocols by ba	aseline conditio	ons (n = 3)		
Baseline condition	Starting dose	Titration		
(+) Diarrhea history	60% TID	Increase by 5 mL TID weekly		
(−) Diarrhea history	60% BID	Increase by 5 mL BID weekly		
(+) Constipation history	50% BID	Increase by 10 mL BID weekly		
(-) Constipation history	10 mL BID	Increase by 10 mL BID weekly		
Goal dose ≤30 mL BID	5 mL BID	Increase by 5 mL BID weekly		
Goal dose ≥40 mL BID	10 mL BID	Increase by 10 mL BID weekly		
Unknown titration protocols (n = 2)				
Starting dose		Titration		
_		Increase by 5–10 mL over 4 steps		
_		Patient-specific and based on tolerability		

Survey question: Please enter your titration protocol(s) for those newly starting trofinetide. Please include dose, frequency of administration, and titration interval BID, twice daily; TID, three times daily



Survey question:

Based on your experience, approximately what percentage of patients are able to achieve their full weight-banded dose after titrating? If you answered less than 100%, for those unable to achieve full dose, approximately what percentage of the full weight-banded dose are the majority able to achieve?

CONCLUSIONS

- Trofinetide dose titration is a common practice among providers at RTT COEs, with several titration strategies available
- Experts reported that dose titration allows for most patients to reach their label dose while experiencing treatment benefit
- Experts evaluate trofinetide efficacy for 4–5 months after the patient achieves their highest tolerable dose after titration
- The clinical approaches reported here correspond with dose adjustments observed in LOTUS, an ongoing real-world evidence study of trofinetide in patients with RTT⁵

REFERENCES

- 1. Neul JL, et al. Ann Neurol. 2010;68(6):944–950.
- 2. Motil KJ, et al. J Pediatr Gastroenterol Nutr. 2012;55(3):292–298.
- 3. DAYBUE (trofinetide) [package insert]. San Diego, CA: Acadia Pharmaceuticals; 2024.
- 4. Neul JL, et al. Nat Med. 2023;29(6):1468–1475.
- 5. Cosand L, et al. Presented at the 2024 Internation Rett Syndrome Foundation Scientific Meeting, June 18–19, 2024, Westminster, CO, USA.

Trofinetide Trial Period

- (Figure 3A)
- dose after titration (**Figure 3B**)



DISCLOSURES

DNL has received funding for clinical trials from Acadia Pharmaceuticals Inc., Anavex Life Sciences Corp., GW Pharmaceuticals, Neurogene, and Rett Syndrome Research Trust, and consultancy fees from Acadia Pharmaceuticals Inc., Neurogene, and Taysha Gene Therapies. **GC** is a consultant to Acadia Pharmaceuticals Inc. **AB** is a consultant to Acadia Pharmaceuticals Inc. **RR** has received funding for clinical trials from Acadia Pharmaceuticals Inc., Anavex Life Sciences Corp., and GW Pharmaceuticals; and funding from the International Rett Syndrome Foundation to support travel expenses. **CWB** is a consultant to Acadia Pharmaceuticals Inc. **EDLR** has received funding for research and consultation from Amicus Therapeutics and Biomarin. **BS** has received funding for clinical trials with Acadia Pharmaceuticals Inc., Grace Science Foundation, Ionis Pharmaceuticals, Marinus Pharmaceuticals, Neuren, Neurogene, and Rett Syndrome Research Trust, as well as funding for consultancy from Neurogene and Taysha Gene Therapies. **AKP** has received funding from the National Institutes of Health, funding for consulting from Acadia Pharmaceuticals Inc., Anavex Life Sciences Corp., AveXis, and GW Pharmaceuticals, is an adviser to the International Rett Syndrome Foundation, and has served as a member of a data safety monitoring board for clinical trials conducted by Taysha Gene Therapies. **ALA** has received funding for clinical trials from Acadia Pharmaceuticals Inc. and Anavex Life Sciences Corp. **CB** is a consultant to Acadia Pharmaceuticals Inc., Harmony Biosciences, and Shionogi Inc. **LC** and **AP** are employees and stakeholders in Acadia Pharmaceuticals Inc.

ACKNOWLEDGMENTS

The study was supported by Acadia Pharmaceuticals Inc. (San Diego, CA, USA). Medical writing support was provided by Juan Sanchez-Cortes, PhD, of Evidence Scientific Solutions, Inc., and funded by Acadia Pharmaceuticals Inc.

• Survey respondents encourage families to allow 4–5 months to determine trofinetide efficacy

• For most, this period of evaluation begins after the patient arrives at their highest tolerable

Survey question:

What is the minimum amount of time that you recommend patients trial trofinetide prior to determining lack of efficacy? Does the minimum amount of time begin when patient starts the medication, achieves full weight-banded dose, or achieves their highest tolerable dose (which may be lower than full weight-banded dose)?



A C A D I A°